

An introduction to Fourier transforms

Isao Sasano

This document is largely based on the reference book [1] with some parts slightly changed.

1 Fourier Integral

Given a function $f(x)$, define $f_L(x)$ as follows.

$$f_L(x) = \begin{cases} f(x) & -L \leq x \leq L \\ 0 & \text{otherwise} \end{cases}$$

Note that

$$f(x) = \lim_{L \rightarrow \infty} f_L(x).$$

Also suppose $f_L(x)$ is represented by a Fourier series

$$f_L(x) = \frac{1}{2}a_0 + \sum_{k=1}^{\infty} (a_k \cos \omega_k x + b_k \sin \omega_k x)$$

on the range $[-L, L]$, where a_0, a_k, b_k, ω_k are given as follows.

$$\begin{aligned} \omega_k &= \frac{k\pi}{L} \\ a_0 &= \frac{1}{L} \int_{-L}^L f_L(x) dx \\ a_k &= \frac{1}{L} \int_{-L}^L f_L(x) \cos \omega_k x dx \\ b_k &= \frac{1}{L} \int_{-L}^L f_L(x) \sin \omega_k x dx \end{aligned}$$

Then $f_L(x)$ is represented as follows.

$$f_L(x) = \frac{1}{2L} \int_{-L}^L f_L(x) dx + \frac{1}{L} \sum_{k=1}^{\infty} \left\{ \cos \omega_k x \int_{-L}^L f_L(x) \cos \omega_k x dx + \sin \omega_k x \int_{-L}^L f_L(x) \sin \omega_k x dx \right\}$$

We now set

$$\Delta\omega = \omega_{k+1} - \omega_k = \frac{(k+1)\pi}{L} - \frac{k\pi}{L} = \frac{\pi}{L}.$$

Since $1/L = \Delta\omega/\pi$ we may rewrite the above equation as follows.

$$\begin{aligned} f_L(x) &= \frac{1}{2L} \int_{-L}^L f_L(x) dx + \frac{\Delta\omega}{\pi} \sum_{k=1}^{\infty} \left\{ \cos \omega_k x \int_{-L}^L f_L(x) \cos \omega_k x dx + \sin \omega_k x \int_{-L}^L f_L(x) \sin \omega_k x dx \right\} \\ &= \frac{1}{2L} \int_{-L}^L f_L(x) dx + \frac{1}{\pi} \sum_{k=1}^{\infty} \left\{ \cos \omega_k x \int_{-L}^L f_L(x) \cos \omega_k x dx + \sin \omega_k x \int_{-L}^L f_L(x) \sin \omega_k x dx \right\} \Delta\omega \\ &= \frac{1}{2L} \int_{-L}^L f_L(x) dx + \frac{1}{\pi} \sum_{k=1}^{\infty} \left\{ \cos k\Delta\omega x \int_{-L}^L f_L(x) \cos k\Delta\omega x dx + \sin k\Delta\omega x \int_{-L}^L f_L(x) \sin k\Delta\omega x dx \right\} \Delta\omega \end{aligned}$$

Now we would like to let $L \rightarrow \infty$, but before that we assume that $f(x)$ is *absolutely integrable* on the x -axis; that is, the following finite limits exist.

$$\lim_{a \rightarrow -\infty} \int_a^0 |f(x)| dx + \lim_{a \rightarrow \infty} \int_0^a |f(x)| dx$$

Note that this is written as $\int_{-\infty}^{\infty} |f(x)| dx$.

We now let $L \rightarrow \infty$, then the first term $\frac{1}{2L} \int_{-L}^L f_L(x) dx$ approaches zero. Also $\Delta\omega = \pi/L$ approaches zero and it seems *plausible* that the infinite series

becomes an integral from 0 to ∞ as follows¹².

$$f(x) = \frac{1}{\pi} \int_0^{\infty} \left\{ \cos \omega x \int_{-\infty}^{\infty} f(x) \cos \omega x dx + \sin \omega x \int_{-\infty}^{\infty} f(x) \sin \omega x dx \right\} d\omega$$

By introducing the functions

$$A(\omega) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(x) \cos \omega x dx \quad (1)$$

and

$$B(\omega) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(x) \sin \omega x dx \quad (2)$$

we can also write the above formula as follows.

$$f(x) = \int_0^{\infty} \{A(\omega) \cos \omega x + B(\omega) \sin \omega x\} d\omega \quad (3)$$

This is called a representation of $f(x)$ by a **Fourier integral**.

The following theorem holds (see p. 513 of the reference book [1]).

Theorem 1 *If $f(x)$ is piecewise continuous in every finite interval and has a right-hand derivative and a left-hand derivative at every point and if $f(x)$ is absolutely integrable, then $f(x)$ can be represented by (3) with A and B given by (1) and (2). At a point where $f(x)$ is discontinuous the value of the Fourier integral equals the average of the left- and right-hand limits of $f(x)$ at that point. In formula,*

$$\int_0^{\infty} \{A(\omega) \cos \omega x + B(\omega) \sin \omega x\} d\omega = \frac{f(x-0) + f(x+0)}{2}.$$

□

¹It just seems plausible and is not the logical consequence, although the equality actually holds similarly to the previous example we have seen in example14-1.pdf, provided that $f(x)$ is piecewise continuous in every finite interval and has a right-hand derivative and a left-hand derivative at every point. In textbooks used in engineering classes, Fourier transform is usually explained as the limit of Fourier series as the period L goes to infinity.

²Notice that I use the same name x for a few variables each of which has different scope. Of course we can rename inner x as other name such as v , which might be a usual way of writing this kind of formula.

Example Calculate the Fourier integral of the following function.

$$f(x) = \begin{cases} 1 & -1 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$

Solution We calculate $A(\omega)$ and $B(\omega)$ as follows.

$$\begin{aligned} A(\omega) &= \frac{1}{\pi} \int_{-\infty}^{\infty} f(x) \cos \omega x dx \\ &= \frac{1}{\pi} \int_{-1}^1 \cos \omega x dx \\ &= \frac{1}{\pi} \left[\frac{\sin \omega x}{\omega} \right]_{-1}^1 \\ &= \frac{2 \sin \omega}{\pi \omega} \end{aligned}$$

$$\begin{aligned} B(\omega) &= \frac{1}{\pi} \int_{-\infty}^{\infty} f(x) \sin \omega x dx \\ &= \frac{1}{\pi} \int_{-1}^1 \sin \omega x dx \\ &= 0 \end{aligned}$$

So we obtain the Fourier integral of $f(x)$ as follows.

$$\frac{2}{\pi} \int_0^{\infty} \frac{\cos \omega x \sin \omega}{\omega} d\omega$$

By Theorem 1 we obtain the following equality.

$$\frac{2}{\pi} \int_0^{\infty} \frac{\cos \omega x \sin \omega}{\omega} d\omega = \begin{cases} 1 & -1 < x < 1 \\ 1/2 & x = -1, 1 \\ 0 & \text{otherwise} \end{cases}$$

We should mention one thing. The integral above is the limit of the formula

$$\frac{2}{\pi} \int_0^a \frac{\cos \omega x \sin \omega}{\omega} d\omega$$

as a goes to infinity. In this integral, there are oscillations near the points $x = -1$ and $x = 1$. The oscillations does not disappear even if a increases, similarly to the Fourier series. This is called the Gibbs phenomenon.

Note By setting $x = 0$ in the Fourier integral of $f(x)$, we obtain the following equality.

$$\frac{2}{\pi} \int_0^{\infty} \frac{\sin \omega}{\omega} d\omega = 1$$

By multiplying the both sides by $\frac{\pi}{2}$ we obtain the following equality.

$$\int_0^{\infty} \frac{\sin \omega}{\omega} d\omega = \frac{\pi}{2}$$

This is called the **Dirichlet integral**. Consider the following (partial) integral (∞ is replaced by a) so called **sine integral**.

$$\text{Si}(a) = \int_0^a \frac{\sin \omega}{\omega} d\omega$$

In the sine integral $\text{Si}(a)$, there are oscillations. The oscillations in the above integral come from the oscillations in the sine integral.

2 Fourier transform

The (real) Fourier integral is

$$f(x) = \int_0^{\infty} \{A(\omega) \cos \omega x + B(\omega) \sin \omega x\} d\omega$$

where A and B are given as follows.

$$\begin{aligned} A(\omega) &= \frac{1}{\pi} \int_{-\infty}^{\infty} f(v) \cos \omega v dv \\ B(\omega) &= \frac{1}{\pi} \int_{-\infty}^{\infty} f(v) \sin \omega v dv \end{aligned}$$

Substituting A and B into the integral, we have

$$\begin{aligned}
f(x) &= \int_0^\infty \left\{ \frac{1}{\pi} \int_{-\infty}^\infty f(v) \cos \omega v dv \cos \omega x + \frac{1}{\pi} \int_{-\infty}^\infty f(v) \sin \omega v dv \sin \omega x \right\} d\omega \\
&= \frac{1}{\pi} \int_0^\infty \left\{ \int_{-\infty}^\infty f(v) \cos \omega v dv \cos \omega x + \int_{-\infty}^\infty f(v) \sin \omega v dv \sin \omega x \right\} d\omega \\
&= \frac{1}{\pi} \int_0^\infty \left\{ \int_{-\infty}^\infty f(v) \cos \omega v \cos \omega x dv + \int_{-\infty}^\infty f(v) \sin \omega v \sin \omega x dv \right\} d\omega \\
&= \frac{1}{\pi} \int_0^\infty \left\{ \int_{-\infty}^\infty f(v) (\cos \omega v \cos \omega x + \sin \omega v \sin \omega x) dv \right\} d\omega \\
&= \frac{1}{\pi} \int_0^\infty \left\{ \int_{-\infty}^\infty f(v) \cos(\omega x - \omega v) dv \right\} d\omega \\
&= \frac{1}{\pi} \int_0^\infty \left\{ \int_{-\infty}^\infty f(v) \cos(\omega(x - v)) dv \right\} d\omega \\
&= \frac{1}{2\pi} \int_{-\infty}^\infty \left\{ \int_{-\infty}^\infty f(v) \cos(\omega(x - v)) dv \right\} d\omega \\
&\quad (\text{since the integral in the brackets is an even function of } \omega)
\end{aligned}$$

The integral of this form with sin instead of cos

$$\frac{1}{2\pi} \int_{-\infty}^\infty \left\{ \int_{-\infty}^\infty f(v) \sin(\omega(x - v)) dv \right\} d\omega$$

is zero since the integral in the brackets is an odd function of ω .

By the Euler formula

$$e^{ix} = \cos x + i \sin x$$

we obtain the following equality.

$$e^{i\omega(x-v)} = \cos(\omega(x - v)) + i \sin(\omega(x - v))$$

By multiplying the both sides by $f(v)$ we obtain the following equality.

$$f(v)e^{i\omega(x-v)} = f(v) \cos(\omega(x - v)) + i f(v) \sin(\omega(x - v))$$

By taking integral with respect to v and ω and multiplying the result by $\frac{1}{2\pi}$

we obtain the following equality.

$$\begin{aligned}
& \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) e^{i\omega(x-v)} dv \right\} d\omega \\
&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) \cos(\omega(x-v)) + i f(v) \sin(\omega(x-v)) dv \right\} d\omega \\
&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) \cos(\omega(x-v)) dv \right\} d\omega \\
&\quad + i \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) \sin(\omega(x-v)) dv \right\} d\omega \\
&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) \cos(\omega(x-v)) dv \right\} d\omega \\
&= f(x)
\end{aligned}$$

So we obtain the following equality.

$$\begin{aligned}
f(x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) e^{i\omega(x-v)} dv \right\} d\omega \\
&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) e^{i\omega x - i\omega v} dv \right\} d\omega \\
&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) e^{i\omega x} e^{-i\omega v} dv \right\} d\omega \\
&= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) e^{-i\omega v} dv \right\} e^{i\omega x} d\omega
\end{aligned}$$

We usually write this as follows³.

$$\begin{aligned}
f(x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) e^{i\omega x} d\omega \\
F(\omega) &= \int_{-\infty}^{\infty} f(x) e^{-i\omega x} dx
\end{aligned}$$

We call $\int_{-\infty}^{\infty} f(x) e^{-i\omega x} dx$ the Fourier transform of $f(x)$ and we call $\frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) e^{i\omega x} d\omega$ the inverse Fourier transform of $F(\omega)$. Note that the

³The constants ($\frac{1}{2\pi}$ and 1) depend on textbooks. They may be $\frac{1}{\sqrt{2\pi}}$ and $\frac{1}{\sqrt{2\pi}}$.

above two equalities are just definitions of transformations and $f(x)$ might not be equal to the inverse Fourier transform of the Fourier transform of $f(x)$ (see Theorem 1).

Example Calculate the Fourier transform of the following function.

$$f(x) = \begin{cases} 1 & -1 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$

Solution

$$\begin{aligned} F(\omega) &= \int_{-1}^1 e^{-i\omega x} dx \\ &= \left[\frac{e^{-i\omega x}}{-i\omega} \right]_{-1}^1 \\ &= \frac{1}{-i\omega} (e^{-i\omega} - e^{i\omega}) \\ &= \frac{1}{-i\omega} (\cos \omega - i \sin \omega - (\cos \omega + i \sin \omega)) \\ &= \frac{1}{-i\omega} (-2i \sin \omega) \\ &= 2 \frac{\sin \omega}{\omega} \end{aligned}$$

References

- [1] Erwin Kreyszig. *Advanced Engineering Mathematics*. John Wiley & Sons Ltd., tenth edition, 2011.